

Search for $B \rightarrow h^{(*)} \nu \bar{\nu}$ Decays at Belle

K.-F. Chen,²⁷ I. Adachi,⁸ H. Aihara,⁴⁵ V. Aulchenko,¹ T. Aushev,^{19,14} S. Bahinipati,³ A. M. Bakich,⁴⁰
V. Balagura,¹⁴ E. Barberio,²² A. Bay,¹⁹ K. Belous,¹³ U. Bitenc,¹⁵ A. Bondar,¹ A. Bozek,²⁸ M. Bračko,^{21,15}
J. Brodzicka,⁸ T. E. Browder,⁷ M.-C. Chang,⁴ P. Chang,²⁷ Y. Chao,²⁷ A. Chen,²⁵ B. G. Cheon,⁶ C.-C. Chiang,²⁷
I.-S. Cho,⁴⁹ Y. Choi,³⁹ Y. K. Choi,³⁹ S. Cole,⁴⁰ M. Danilov,¹⁴ M. Dash,⁴⁸ A. Drutskoy,³ S. Eidelman,¹ S. Fratina,¹⁵
N. Gabyshev,¹ B. Golob,^{20,15} H. Ha,¹⁷ J. Haba,⁸ T. Hara,³³ K. Hayasaka,²³ M. Hazumi,⁸ D. Heffernan,³³
T. Hokuue,²³ Y. Hoshi,⁴³ W.-S. Hou,²⁷ Y. B. Hsiung,²⁷ H. J. Hyun,¹⁸ T. Iijima,²³ K. Ikado,²³ K. Inami,²³
A. Ishikawa,⁴⁵ H. Ishino,⁴⁶ R. Itoh,⁸ M. Iwasaki,⁴⁵ Y. Iwasaki,⁸ N. J. Joshi,⁴¹ S. Kajiwarra,³³ J. H. Kang,⁴⁹
N. Katayama,⁸ H. Kawai,² T. Kawasaki,³⁰ H. Kichimi,⁸ Y. J. Kim,⁵ K. Kinoshita,³ S. Korpar,^{21,15}
P. Križan,^{20,15} P. Krokovny,⁸ R. Kumar,³⁴ C. C. Kuo,²⁵ A. Kuzmin,¹ Y.-J. Kwon,⁴⁹ J. S. Lee,³⁹ S. E. Lee,³⁸
T. Lesiak,²⁸ S.-W. Lin,²⁷ Y. Liu,⁵ D. Liventsev,¹⁴ F. Mandl,¹² D. Marlow,³⁵ A. Matyja,²⁸ S. McOnie,⁴⁰
T. Medvedeva,¹⁴ K. Miyabayashi,²⁴ H. Miyake,³³ H. Miyata,³⁰ Y. Miyazaki,²³ R. Mizuk,¹⁴ Y. Nagasaka,⁹
I. Nakamura,⁸ M. Nakao,⁸ S. Nishida,⁸ S. Ogawa,⁴² T. Ohshima,²³ S. Okuno,¹⁶ S. L. Olsen,⁷ H. Ozaki,⁸
P. Pakhlov,¹⁴ G. Pakhlova,¹⁴ H. Park,¹⁸ K. S. Park,³⁹ R. Pestotnik,¹⁵ L. E. Piilonen,⁴⁸ Y. Sakai,⁸ O. Schneider,¹⁹
J. Schümann,⁸ C. Schwanda,¹² A. J. Schwartz,³ R. Seidl,^{10,36} K. Senyo,²³ M. E. Sevier,²² M. Shapkin,¹³
C. P. Shen,¹¹ H. Shibuya,⁴² S. Shinomiya,³³ J.-G. Shiu,²⁷ B. Shwartz,¹ J. B. Singh,³⁴ A. Sokolov,¹³ A. Somov,³
S. Stanič,³¹ M. Starič,¹⁵ K. Sumisawa,⁸ T. Sumiyoshi,⁴⁷ O. Tajima,⁸ F. Takasaki,⁸ N. Tamura,³⁰ M. Tanaka,⁸
G. N. Taylor,²² Y. Teramoto,³² I. Tikhomirov,¹⁴ K. Trabelsi,⁸ S. Uehara,⁸ K. Ueno,²⁷ T. Uglov,¹⁴ Y. Unno,⁶
S. Uno,⁸ P. Urquijo,²² Y. Usov,¹ G. Varner,⁷ K. E. Varvell,⁴⁰ K. Vervink,¹⁹ S. Villa,¹⁹ A. Vinokurova,¹
C. C. Wang,²⁷ C. H. Wang,²⁶ M.-Z. Wang,²⁷ P. Wang,¹¹ Y. Watanabe,¹⁶ R. Wedd,²² E. Won,¹⁷ B. D. Yabsley,⁴⁰
A. Yamaguchi,⁴⁴ Y. Yamashita,²⁹ M. Yamauchi,⁸ C. C. Zhang,¹¹ Z. P. Zhang,³⁷ V. Zhilich,¹ and A. Zupanc¹⁵

(The Belle Collaboration)

¹*Budker Institute of Nuclear Physics, Novosibirsk*²*Chiba University, Chiba*³*University of Cincinnati, Cincinnati, Ohio 45221*⁴*Department of Physics, Fu Jen Catholic University, Taipei*⁵*The Graduate University for Advanced Studies, Hayama*⁶*Hanyang University, Seoul*⁷*University of Hawaii, Honolulu, Hawaii 96822*⁸*High Energy Accelerator Research Organization (KEK), Tsukuba*⁹*Hiroshima Institute of Technology, Hiroshima*¹⁰*University of Illinois at Urbana-Champaign, Urbana, Illinois 61801*¹¹*Institute of High Energy Physics, Chinese Academy of Sciences, Beijing*¹²*Institute of High Energy Physics, Vienna*¹³*Institute of High Energy Physics, Protvino*¹⁴*Institute for Theoretical and Experimental Physics, Moscow*¹⁵*J. Stefan Institute, Ljubljana*¹⁶*Kanagawa University, Yokohama*¹⁷*Korea University, Seoul*¹⁸*Kyungpook National University, Taegu*¹⁹*Swiss Federal Institute of Technology of Lausanne, EPFL, Lausanne*²⁰*University of Ljubljana, Ljubljana*²¹*University of Maribor, Maribor*²²*University of Melbourne, School of Physics, Victoria 3010*²³*Nagoya University, Nagoya*²⁴*Nara Women's University, Nara*²⁵*National Central University, Chung-li*²⁶*National United University, Miao Li*²⁷*Department of Physics, National Taiwan University, Taipei*²⁸*H. Niewodniczanski Institute of Nuclear Physics, Krakow*²⁹*Nippon Dental University, Niigata*³⁰*Niigata University, Niigata*³¹*University of Nova Gorica, Nova Gorica*³²*Osaka City University, Osaka*

- ³³Osaka University, Osaka
³⁴Panjab University, Chandigarh
³⁵Princeton University, Princeton, New Jersey 08544
³⁶RIKEN BNL Research Center, Upton, New York 11973
³⁷University of Science and Technology of China, Hefei
³⁸Seoul National University, Seoul
³⁹Sungkyunkwan University, Suwon
⁴⁰University of Sydney, Sydney, New South Wales
⁴¹Tata Institute of Fundamental Research, Mumbai
⁴²Toho University, Funabashi
⁴³Tohoku Gakuin University, Tagajo
⁴⁴Tohoku University, Sendai
⁴⁵Department of Physics, University of Tokyo, Tokyo
⁴⁶Tokyo Institute of Technology, Tokyo
⁴⁷Tokyo Metropolitan University, Tokyo
⁴⁸Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061
⁴⁹Yonsei University, Seoul

We present a search for the rare decays $B \rightarrow h^{(*)}\nu\bar{\nu}$, where $h^{(*)}$ stands for a light meson. A data sample of 535 million $B\bar{B}$ pairs collected with the Belle detector at the KEKB e^+e^- collider is used. Signal candidates are required to have an accompanying B meson fully reconstructed in a hadronic mode and signal-side particles consistent with a single $h^{(*)}$ meson. No significant signal is observed and we set upper limits on the branching fractions at 90% confidence level. The limits on $B^0 \rightarrow K^{*0}\nu\bar{\nu}$ and $B^+ \rightarrow K^+\nu\bar{\nu}$ decays are more stringent than the previous constraints, while the first searches for $B^0 \rightarrow K^0\nu\bar{\nu}$, $\pi^0\nu\bar{\nu}$, $\rho^0\nu\bar{\nu}$, $\phi\nu\bar{\nu}$ and $B^+ \rightarrow K^{*+}\nu\bar{\nu}$, $\rho^+\nu\bar{\nu}$ are reported.

PACS numbers: 13.25.Hw, 14.40.Nd

The decays $B \rightarrow K^{(*)}\nu\bar{\nu}$ proceed through the flavor-changing neutral-current process $b \rightarrow s\nu\bar{\nu}$, which is sensitive to physics beyond the Standard Model (SM). The dominant SM diagrams are shown in Fig. 1. Similarly, the decays $B \rightarrow (\pi, \rho)\nu\bar{\nu}$ proceed through $b \rightarrow d\nu\bar{\nu}$ processes. The SM branching fractions are estimated to be 1.3×10^{-5} and 4×10^{-6} for $B \rightarrow K^*\nu\bar{\nu}$ and $B \rightarrow K\nu\bar{\nu}$ decays [1], respectively, and are expected to be much lower for other modes. Theoretical calculation of the decay amplitudes for $B \rightarrow h^{(*)}\nu\bar{\nu}$ is particularly reliable, owing to the absence of long-distance interactions that affect charged-lepton channels $B \rightarrow h^{(*)}l^+l^-$. New physics such as SUSY particles or a possible fourth generation could potentially contribute to the penguin loop or box diagram and enhance the branching fractions [1]. Reference [2] also discusses the possibility of discovering light dark matter in $b \rightarrow s$ transitions with large missing momentum.




FIG. 1: The quark-level diagrams for $B \rightarrow K^*\nu\bar{\nu}$ decays.

Experimental measurements [3] of the $b \rightarrow s$ transitions with two charged leptons are in good agreement with SM calculations [1]. Further investigation of the forward-backward asymmetry in $B \rightarrow K^*l^+l^-$ [4] is

consistent with the SM although the statistics are still limited. Due to the challenge of cleanly detecting rare modes with two final-state neutrinos, only a few studies of $K^{(*)}\nu\bar{\nu}$ have been carried out to date [5, 6, 7]; there is only one examination of the corresponding $b \rightarrow d$ transitions [7]. In this paper, we report our first search for the decays $B \rightarrow h^{(*)}\nu\bar{\nu}$ ($h^{(*)}$ stands for K^+ , K_S^0 , K^{*0} , K^{*+} , π^+ , π^0 , ρ^0 , ρ^+ , and ϕ) using a 492 fb^{-1} data sample recorded at the $\Upsilon(4S)$ resonance, corresponding to 535×10^6 B -meson pairs. Charge-conjugate decays are implied throughout this paper.

The Belle detector is a large-solid-angle magnetic spectrometer located at the KEKB collider [8], and consists of a silicon vertex detector (SVD), a 50-layer central drift chamber (CDC), an array of aerogel threshold Cherenkov counters (ACC), a barrel-like arrangement of time-of-flight scintillation counters (TOF), and an electromagnetic calorimeter comprised of CsI(Tl) crystals (ECL) located inside a superconducting solenoid that provides a 1.5 T magnetic field. An iron flux-return located outside the coil is instrumented to detect K_L^0 mesons and to identify muons (KLM). The detector is described in detail elsewhere [9].

Candidate $e^+e^- \rightarrow \Upsilon(4S) \rightarrow B\bar{B}$ events are characterized by a fully-reconstructed tag-side B meson (B_{tag}). The remaining particles are assumed to be products of the signal-side B meson (B_{sig}). The B_{tag} candidates are reconstructed in one of the following modes: $B^0 \rightarrow D^{(*)-}\pi^+$, $D^{(*)-}\rho^+$, $D^{(*)-}a_1^+$, $D^{(*)-}D_s^{(*)+}$, $B^+ \rightarrow \bar{D}^{(*)0}\pi^+$, $\bar{D}^{(*)0}\rho^+$, $\bar{D}^{(*)0}a_1^+$, and $\bar{D}^{(*)0}D_s^{(*)+}$.

The D^- mesons are reconstructed as $D^- \rightarrow K_S^0 \pi^-$, $K_S^0 \pi^- \pi^0$, $K_S^0 \pi^- \pi^+ \pi^-$, $K^+ \pi^- \pi^-$, and $K^+ \pi^- \pi^- \pi^0$. The following decay channels are included for $\bar{D}^0$ mesons: $\bar{D}^0 \rightarrow K^+ \pi^-$, $K^+ \pi^- \pi^0$, $K^+ \pi^- \pi^+ \pi^-$, $K_S^0 \pi^0$, $K_S^0 \pi^- \pi^+$, $K_S^0 \pi^- \pi^+ \pi^0$ and $K^- K^+$. The D^{*-} ($\bar{D}^{*0}$) mesons are reconstructed as $\bar{D}^0 \pi^-$ ($\bar{D}^0 \pi^0$ and $\bar{D}^0 \gamma$). Furthermore, $D_s^{*+} \rightarrow D_s^+ \gamma$, $D_s^+ \rightarrow K_S^0 K^+$ and $K^+ K^- \pi^+$ decays are reconstructed. B_{tag} candidates are selected using the beam-energy constrained mass $M_{\text{bc}} \equiv \sqrt{E_{\text{beam}}^2 - p_B^2}$ and the energy difference $\Delta E \equiv E_B - E_{\text{beam}}$, where E_B and p_B are the reconstructed energy and momentum of the B_{tag} candidate in the $\Upsilon(4S)$ center-of-mass (CM) frame, and E_{beam} is the beam energy in this frame.

We require B_{tag} candidates satisfy the requirements $M_{\text{bc}} > 5.27 \text{ GeV}/c^2$ and $-80 \text{ MeV} < \Delta E < 60 \text{ MeV}$. If there are multiple B_{tag} candidates in an event, the candidate with the smallest χ^2 based on the deviations from the nominal values of ΔE , the D meson mass, and the mass difference between the D^* and the D (for candidates with a D^* in the final state) is chosen. We reconstruct 7.88×10^5 and 4.91×10^5 charged and neutral B mesons, respectively.

The particles in the event not associated with the B_{tag} meson are used to reconstruct a $B_{\text{sig}} \rightarrow h^{(*)} \nu \bar{\nu}$ candidate. Prompt charged tracks are required to have a maximum distance to the interaction point (IP) of 5 cm in the beam direction (z), of 2 cm in the transverse plane (r - ϕ), and a minimum momentum of 0.1 GeV/ c in the transverse plane. We reconstruct $K^\pm$ ($\pi^\pm$) candidates from charged tracks having a kaon likelihood greater than 0.6 (less than 0.4) with an efficiency of 84–91% (87–92%). The kaon likelihood is defined by $\mathcal{R}_K \equiv \mathcal{L}_K / (\mathcal{L}_K + \mathcal{L}_\pi)$, where $\mathcal{L}_K$ ($\mathcal{L}_\pi$) denotes a combined likelihood measurement from the ACC, the TOF, and a dE/dx from the CDC for the $K^\pm$ ($\pi^\pm$) tracks. Pairs of oppositely charged tracks are used to reconstruct $K_S^0 \rightarrow \pi^+ \pi^-$ decays, with an invariant mass that is within $\pm 15 \text{ MeV}/c^2$ ($> 5 \sigma$) from the nominal K_S^0 meson mass. The $\pi^+ \pi^-$ vertex is required to be displaced from the IP by a minimum distance of 0.22 cm. The direction of the pion pair momentum in the transverse plane must agree with the direction defined by the IP and the vertex displacement within 0.03 rad. For $\pi^0 \rightarrow \gamma \gamma$, a minimum photon energy of 50 MeV is required and the $\gamma \gamma$ invariant mass must be within $\pm 16 \text{ MeV}/c^2$ ($\sim 2.5 \sigma$) of the nominal π^0 mass.

The decays $B_{\text{sig}} \rightarrow K^+ \nu \bar{\nu}$, $\pi^+ \nu \bar{\nu}$, $K_S^0 \nu \bar{\nu}$, and $\pi^0 \nu \bar{\nu}$ are reconstructed from single K^+ , π^+ , K_S^0 , and π^0 candidates, respectively. The $B^0 \rightarrow K^{*0} \nu \bar{\nu}$ candidate is reconstructed from a charged pion and an oppositely charged kaon, while $B^+ \rightarrow K^{*+} \nu \bar{\nu}$ decays are reconstructed from a K_S^0 candidate and a charged pion, or a charged kaon and a π^0 candidate. The reconstructed mass of the K^{*0} (K^{*+}) candidate should be within a $\pm 75 \text{ MeV}/c^2$ window around the nominal K^{*0} (K^{*+}) mass. Furthermore, pairs of charged pions with opposite charge are used to form $B^0 \rightarrow \rho^0 \nu \bar{\nu}$ candidates where the $\pi^+ \pi^-$ invariant

mass should be within $\pm 150 \text{ MeV}/c^2$ from the nominal ρ^0 mass. For $B^+ \rightarrow \rho^+ \nu \bar{\nu}$, a charged pion and a π^0 candidate are used, and a $\pm 150 \text{ MeV}/c^2$ mass window is required. A ϕ meson is formed from a $K^+ K^-$ pair with a reconstructed mass within $\pm 10 \text{ MeV}/c^2$ ($\sim 2 \sigma$) from the nominal ϕ mass.

No additional charged tracks or π^0 candidates are allowed in the event. We select B_{sig} candidates using the variable $E_{\text{ECL}} \equiv E_{\text{tot}} - E_{\text{rec}}$, where E_{tot} and E_{rec} are the total visible energy measured by the ECL detector and the measured energy of reconstructed objects including the B_{tag} and the signal side $h^{(*)}$ candidate, respectively. A minimum threshold of 50 (100, 150) MeV on the cluster energy is applied for the barrel (forward endcap, backward endcap) region of the ECL detector. The decays $B \rightarrow D^* \ell \nu$ are examined as control samples; the observed E_{ECL} distributions are found to be in good agreement with Monte Carlo (MC) simulations [10]. The signal region is defined by $E_{\text{ECL}} < 0.3 \text{ GeV}$ while the sideband region is given by $0.45 \text{ GeV} < E_{\text{ECL}} < 1.5 \text{ GeV}$.

The dominant background source is $B\bar{B}$ decays involving a $b \rightarrow c$ transition. A lower bound of 1.6 GeV/ c on P^* , the momentum of the $h^{(*)}$ candidate in the B_{sig} rest frame, suppresses this background, while an upper bound of 2.5 GeV/ c rejects the contributions from radiative two-body modes such as $B \rightarrow K^* \gamma$. The P^* requirement is removed for ϕ candidates due to the lack of theoretical calculations for $B_d \rightarrow \phi$ form factors. Furthermore, the cosine of the angle between the missing momentum in the laboratory frame and the beam is required to lie between -0.86 and 0.95 . The missing momentum is calculated using the momenta of the reconstructed B_{tag} and $h^{(*)}$ candidates. These criteria suppress backgrounds with particles produced along the beam pipe. Other background sources, such as $e^+ e^- \rightarrow q \bar{q}$ ($q = u, d, c, s$) continuum background and rare B decays involving $b \rightarrow u$, $b \rightarrow s$, or $b \rightarrow d$ processes, are found to be small.

The reconstructed E_{ECL} distributions are shown in Fig. 2. The E_{ECL} distributions of background are estimated with MC simulations; in particular, a large $b \rightarrow c$ MC sample corresponding to ten times the data luminosity is introduced with a preselection on the generator information. The background E_{ECL} distributions are normalized by the number of events in the sideband region. None of the signal modes has a significant signal. Including the effects of both statistical and systematic uncertainties, an extension of the Feldman-Cousins method [11, 12] is used to calculate the upper limits on the branching fractions. The observed number of events in the signal box and sideband region, expected background contributions in the signal box, reconstruction efficiencies, and the obtained upper limits at 90% confidence level (CL) are shown in Table I. The reconstruction efficiencies are estimated with MC simulations using the $B \rightarrow h^{(*)}$ form factors from Ref. [13]. The $B^0 \rightarrow \phi \nu \bar{\nu}$ MC samples are generated with the $B \rightarrow K^*$ form factors.




FIG. 2: The E_{ECL} distributions for $B \rightarrow h^{(*)}\nu\bar{\nu}$ decays. The shaded histograms show the background distributions from MC simulations and are normalized to sideband data. The open histograms show the SM expected signal distributions for $B \rightarrow K^{(*)}\nu\bar{\nu}$ decays multiplied by a factor of 20 for the comparison. The vertical dashed lines show the upper bound (left) of the signal box and the lower bound (right) of the sideband region.

TABLE I: A summary of the number of observed events in the signal box (N_{obs}) and sideband regions (N_{side}), expected background yields (N_b) in the signal box, reconstruction efficiencies including both B_{tag} and B_{sig} (ϵ), and the upper limits (U.L.) on the branching fractions at 90% CL.

Mode	N_{obs}	N_{side}	N_b	$\epsilon(\times 10^{-5})$	U.L.
$K^{*0}\nu\bar{\nu}$	7	16	4.2 ± 1.4	5.1 ± 0.3	$< 3.4 \times 10^{-4}$
$K^{*+}\nu\bar{\nu}$	4	18	5.6 ± 1.8	5.8 ± 0.7	$< 1.4 \times 10^{-4}$
$\rightarrow K_S^0\pi^+$	1	7	2.3 ± 1.2	2.8 ± 0.3	
$\rightarrow K^+\pi^0$	3	11	3.3 ± 1.4	3.0 ± 0.4	
$K^+\nu\bar{\nu}$	10	60	20.0 ± 4.0	26.7 ± 2.9	$< 1.4 \times 10^{-5}$
$K^0\nu\bar{\nu}$	2	8	2.0 ± 0.9	5.0 ± 0.3	$< 1.6 \times 10^{-4}$
$\pi^+\nu\bar{\nu}$	33	149	25.9 ± 3.9	24.2 ± 2.6	$< 1.7 \times 10^{-4}$
$\pi^0\nu\bar{\nu}$	11	15	3.8 ± 1.3	12.8 ± 0.8	$< 2.2 \times 10^{-4}$
$\rho^0\nu\bar{\nu}$	21	46	11.5 ± 2.3	8.4 ± 0.5	$< 4.4 \times 10^{-4}$
$\rho^+\nu\bar{\nu}$	15	66	17.8 ± 3.2	8.5 ± 1.1	$< 1.5 \times 10^{-4}$
$\phi\nu\bar{\nu}$	1	9	1.9 ± 0.9	9.6 ± 1.4	$< 5.8 \times 10^{-5}$

The uncertainties associated with the background yields are dominated by the data sideband statistics and MC statistics, and are summarized in Table II. The possible disagreement in the E_{ECL} distributions between data and MC is checked using wrong-flavor combinatorial

events, and an uncertainty of 0.1–2.0 events is included. Possible backgrounds from rare B decays are examined using a large MC sample corresponding to 50 times the data luminosity. We change the relative normalizations of rare B components by $\pm 50\%$, and the variation in the background yield (0.1–1.8 events) is included as a systematic uncertainty.

Various sources of uncertainties are considered for the signal normalization and are summarized in Table III. The uncertainties in B_{tag} reconstruction (2.0% for B^0 and 9.9% for B^+) are estimated by comparing the yields of data and MC from the B_{tag} candidates. Systematic uncertainty arising from the track and π^0 rejection is studied using $B \rightarrow D^{(*)}\ell\nu$ decays, and an error of 2.7% is assigned. The uncertainties in the efficiencies for detecting a K_S^0 or π^0 from B_{sig} are estimated to be 4.9% and 4.0%, respectively. We also vary the $B \rightarrow h^{(*)}$ form factors used in the signal MC generation according to the uncertainties given by the Ref. [13], and an uncertainty of 0.4–3.7% on the reconstruction efficiency is included. A larger uncertainty of 13%, which is estimated from the difference between the default decay model and a generic three-body phase-space model, is introduced for $B^0 \rightarrow \phi\nu\bar{\nu}$ decays. Furthermore, the following uncertainties are also considered: the number of $B\bar{B}$ events (1.3%), tracking efficiency (1.0–2.2%), particle identifica-

tion (0.5–2.0%), $h^{(*)}$ mass selection (0.8–2.3%), and the $\phi \rightarrow K^+ K^-$ branching fraction (1.2%).

In conclusion, we have performed a search for $B \rightarrow h^{(*)} \nu \bar{\nu}$ decays with a fully reconstructed B tagging method on a data sample of 535×10^6 $B\bar{B}$ pairs collected at the $\Upsilon(4S)$ resonance with the Belle detector. No significant signal is observed and we set upper limits on the branching fractions at 90% CL. The limits obtained for $B^0 \rightarrow K^{*0} \nu \bar{\nu}$ and $B^+ \rightarrow K^+ \nu \bar{\nu}$ decays are more stringent than the previous constraints from DELPHI [5] and BaBar [7]. The first searches for $B^0 \rightarrow K^0 \nu \bar{\nu}$, $\pi^0 \nu \bar{\nu}$, $\rho^0 \nu \bar{\nu}$, $\phi \nu \bar{\nu}$, and $B^+ \rightarrow K^{*+} \nu \bar{\nu}$, $\rho^+ \nu \bar{\nu}$ are carried out, and upper limits on the branching fraction of order 10^{-4} are obtained. The limit on $B^+ \rightarrow \pi^+ \nu \bar{\nu}$ is less restrictive than BaBar's result [7] due to a larger number of observed events in the signal box. The results on $B \rightarrow K^{(*)} \nu \bar{\nu}$ reported here are one order of magnitude above the predictions of Buchalla *et al.* [1] and hence still allow room for substantial non-SM contributions. A higher luminosity B -factory experiment is required to probe the SM predictions for the branching fractions.

We thank the KEKB group for excellent operation of the accelerator, the KEK cryogenics group for efficient solenoid operations, and the KEK computer group and the NII for valuable computing and Super-SINET network support. We acknowledge support from MEXT and JSPS (Japan); ARC and DEST (Australia); NSFC and KIP of CAS (China); DST (India); MOEHRD, KOSEF, KRF and SBS Foundation (Korea); KBN (Poland); MES and RFAAE (Russia); ARRS (Slovenia); SNSF (Switzerland); NSC and MOE (Taiwan); and DOE (USA).

-
- [1] G. Buchalla, G. Hiller and G. Isidori, Phys. Rev. D **63**, 014015 (2000).
 - [2] C. Bird, P. Jackson, R. Kowalewski and M. Pospelov, Phys. Rev. Lett. **93**, 201803 (2004).
 - [3] K. Abe *et al.* (Belle Collaboration), Phys. Rev. Lett. **88**, 021801 (2002); A. Ishikawa *et al.* (Belle Collaboration), Phys. Rev. Lett. **91**, 261601 (2003); M. Iwasaki *et al.* (Belle Collaboration), Phys. Rev. D **72**, 092005 (2005); B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **73**, 092001 (2006); B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. Lett. **93**, 081802 (2004).
 - [4] A. Ishikawa *et al.* (Belle Collaboration), Phys. Rev. Lett. **96**, 251801 (2006).
 - [5] W. Adam *et al.* (DELPHI Collaboration), Z. Phys. C **72**, 207 (1996).
 - [6] T. E. Browder *et al.* (CLEO Collaboration), Phys. Rev. Lett. **86**, 2950 (2001).
 - [7] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. Lett. **94**, 101801 (2005).
 - [8] S. Kurokawa and E. Kikutani, Nucl. Instrum. Methods Phys. Res., Sect. A **499**, 1 (2003).
 - [9] A. Abashian *et al.* (Belle Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 117 (2002).
 - [10] K. Ikado *et al.*, Phys. Rev. Lett. **97**, 251802 (2006).
 - [11] G. J. Feldman and R. D. Cousins, Phys. Rev. D **57**, 3873 (1998).
 - [12] J. Conrad, O. Botner, A. Hallgren and C. Perez de los Heros, Phys. Rev. D **67**, 012002 (2003).
 - [13] P. Ball and R. Zwicky, Phys. Rev. D **71**, 014015 (2005); Phys. Rev. D **71**, 014029 (2005).

TABLE II: Summary of the uncertainties associated with the background yields (in the number of events).

Uncertainties	$K^{*0}\nu\bar{\nu}$	$K^{*+}\nu\bar{\nu}$	$K^+\nu\bar{\nu}$	$K^0\nu\bar{\nu}$	$\pi^+\nu\bar{\nu}$	$\pi^0\nu\bar{\nu}$	$\rho^0\nu\bar{\nu}$	$\rho^+\nu\bar{\nu}$	$\phi\nu\bar{\nu}$
Sideband statistics	1.0	1.3	2.6	0.7	2.1	1.0	1.7	2.2	0.6
MC statistics	0.9	1.2	2.6	0.6	2.0	0.8	1.3	1.6	0.6
MC/data difference	0.3	0.3	1.5	0.2	2.0	0.3	0.9	1.4	0.1
Rare B	0.1	0.3	0.2	0.2	1.8	0.1	0.1	0.9	0.3
Total	1.4	1.8	4.0	0.9	3.9	1.3	2.3	3.2	0.9

TABLE III: Summary of the relative uncertainties for signal normalization (in %).

Source	$K^{*0}\nu\bar{\nu}$	$K^{*+}\nu\bar{\nu}$	$K^+\nu\bar{\nu}$	$K^0\nu\bar{\nu}$	$\pi^+\nu\bar{\nu}$	$\pi^0\nu\bar{\nu}$	$\rho^0\nu\bar{\nu}$	$\rho^+\nu\bar{\nu}$	$\phi\nu\bar{\nu}$
$N(B\bar{B})$	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3
Tracking efficiency	2.1	1.1	1.0	-	1.0	-	2.2	1.1	2.0
K_S^0/π^0 reconstruction	-	4.4	-	4.9	-	4.0	-	4.0	-
Sub branching fraction	-	-	-	-	-	-	-	-	1.2
Particle identification	1.3	0.7	0.7	-	0.5	-	1.0	0.5	2.0
MC statistics	3.5	2.4	1.9	3.2	2.0	2.8	3.3	3.4	2.3
Mass selection	0.8	2.3	-	-	-	-	1.1	2.6	2.0
B_{tag} reconstruction	2.0	9.9	9.9	2.0	9.9	2.0	2.0	9.9	2.0
Track/ π^0 rejection	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7
$B \rightarrow h^{(*)}$ form factor	1.6	1.3	2.6	0.4	3.0	1.0	1.7	3.7	13.0
Total	5.9	11.8	10.9	6.9	11.0	6.1	5.8	12.5	14.2